



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Exploiting body mechanics to produce exploratory behaviour.

Citation for published version:

Ross, D, Lagogiannis, K & Webb, B 2016, 'Exploiting body mechanics to produce exploratory behaviour.', Janelia Behavioral Neurogenetics of Drosophila Larva 2016, United Kingdom, 23/10/16 - 26/10/16.

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record


General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Exploiting body mechanics to produce exploratory behaviour.

Dylan Ross, Konstantinos Lagogiannis, Barbara Webb

Institute of Perception, Action, and Behaviour, School of Informatics, University of Edinburgh, 10 Crichton St, Edinburgh EH8 9AB, UK.

dylrxs@gmail.com



1. Introduction

Body mechanics and sensory feedback are central to the production of rectilinear peristaltic locomotion in the *Drosophila* larva [1; 2; 3; 4]. However, their importance for this animal's lateral motion and substrate exploration is unclear.

We present a model of planar larval mechanics coupled to a simple, reflexive nervous system. Our conservative mechanics produces axial compression waves and chaotic transverse bending. On addition of damping and driving forces, these motions are harnessed to produce forward and backward locomotion as well as deterministic chaotic exploratory behaviour. This is particularly surprising given that our neural model neither explicitly encodes nor controls the direction of movement (either forward/backward, or left/right).

2. Model construction




Figure 1: a mechanical model of midline motion in the plane

- The larval midline is modelled as a set of discrete point masses.
- Elastic energy is stored in axial compression/expansion and transverse bending.
- Energy is lost to viscous friction during motion.
- Masses interact with the substrate via Coulomb sliding friction.
- Head and tail are coupled via a total length constraint.




Figure 2: segmental reflexes interacting via mutual inhibition

- Neurons and muscles are binary (ON/OFF) and have no intrinsic dynamics.
- A reflex loop produces muscle tension within a segment when it is shortening.
- When a reflex is active, it inhibits distant neighbours.

3. Small-amplitude behaviour




Figure 3: modal analysis of conservative mechanics




Figure 4: emergence of limit cycles for forward and backward locomotion

4. Large-amplitude behaviour




Figure 5: conservative head motions become chaotic at large amplitudes




Figure 6: deterministic chaotic exploration

5. Conclusions

- Conservative body mechanics produces motions (axial travelling waves, chaotic transverse bending) suitable for substrate exploration.
- This allows stable forward and backward locomotion as well as turning/exploration to be generated by a single, simple reflex circuit.
- Intrinsic pattern generation and explicit encoding and control of the direction of travel are, in principle, unnecessary for successful exploration.

References

- [1] Song, W., Maika, O., Jan, L.Y., Jan, Y.N.: Peripheral multidendritic sensory neurons are necessary for rhythmic locomotion behaviour in *Drosophila* larvae *PNAS*, 104 (12): 5199–5204, (2007).
- [2] Gjorgjieva J., Jimena, B., Evers, J.F., Eglén, S.: Neural circuits for peristaltic wave propagation in crawling *Drosophila* larvae: analysis and modeling *Frontiers in Computational Neuroscience*, 7:1–24, (2013).
- [3] Ross, D., Lagogiannis, K., Webb, B.: A model of larval biomechanics reveals exploitable passive properties for efficient locomotion. *Lecture Notes in Computer Science*, 9222:1–12, (2015).
- [4] Pehlevan, C., Paoletti, P., Mahadevan, L.: Integrative neuromechanics of crawling in *D. melanogaster* larvae. *eLife*, 5:e11031, (2016).